

On certain canonical diffeomorphisms in symplectic and Poisson geometry

K. C. H. Mackenzie

ABSTRACT. Associated with the canonical symplectic structure on a cotangent bundle T^*M is the diffeomorphism $\#: T^*(T^*M) \rightarrow T(T^*M)$. This and the Tulczyjew diffeomorphism $T(T^*M) \rightarrow T^*(TM)$ may be derived from the canonical involution $T(TM) \rightarrow T(TM)$ by suitable dualizations. We show that the constructions which yield these maps extend very generally to the double Lie algebroids of double Lie groupoids, where they play a crucial role in the relations between double Lie algebroids and Lie bialgebroids.

There have been several talks this meeting about notions of double for Lie bialgebroids. Some of these have derived from the 1997 construction of Liu Zhang–Ju, Alan Weinstein and Xu Ping [4] in which they introduced the notion of Courant algebroid, and some have involved elements of super mathematics. I very much hope that before long there will be a clear account of the relations between these various approaches and even a unification of them.

There is another approach to the question of doubles, which was not at first related to Lie bialgebroids, but arose out of broad considerations of what may be called “second-order geometry”. It is not possible to describe this approach from scratch in an hour, but it is appropriate at this conference to indicate the broad features of it. This talk therefore takes a slice through the papers [5], [7], [6], [8], transverse to their chronological sequence and provides an alternative route to approach them. Some aspects of §2 and §4 are new.

I am very grateful to Ted Voronov and Mike Prest for the splendid opportunities and good fellowship which the Workshop provided, and to the London Mathematical Society for its support. I also wish to thank Yvette Kosmann–Schwarzbach for her comments on an earlier version.

1.

The three canonical diffeomorphisms with which the paper begins are associated with iterated tangent and cotangent bundles:

- (1) The canonical involution on an iterated tangent bundle

$$J: T^2M \rightarrow T^2M,$$

2000 *Mathematics Subject Classification.* Primary 58H01. Secondary 17B66, 18D05, 22A22, 58F05.

which can be loosely described as ‘interchanging the order of differentiation’.

Regard elements of $T^2M = T(TM)$ as represented by equivalence classes of squares $\varphi: I^2 \rightarrow M$. Thus $\varphi = \varphi(s, t)$ differentiates to a curve of tangent vectors $X_t = \frac{\partial \varphi}{\partial s}(0, t)$ and X_t in turn differentiates to $\xi = \frac{d}{dt}X_t|_0 \in T_{X_0}(TM)$. Now J is induced by the map $\varphi \mapsto \varphi \circ \sigma$ where $\sigma(s, t) = (t, s)$.

(2) The Poisson anchor for the canonical symplectic structure on a cotangent bundle T^*M :

$$\pi_{d\theta}^\#: T^*(T^*M) \rightarrow T(T^*M).$$

Any Poisson structure on a manifold P induces a map $\#: T^*P \rightarrow TP$ which sends an exact differential δf to the Hamiltonian vector field X_f . This map serves as the anchor of the Lie algebroid structure defined by the bracket of 1-forms. When P is the cotangent bundle $c: T^*M \rightarrow M$, the canonical symplectic structure is $\omega = \delta\theta$ where θ is the 1-form which associates to $\eta \in T_\alpha(T^*M)$ the pairing of α with $T(c)(\eta) \in TM$.

(3) The canonical diffeomorphism

$$\Theta: T(T^*M) \rightarrow T^*(TM)$$

defined intrinsically by Tulczyjew in [12, p.97] and locally in, for example, [1, p.424]. We call this the *Tulczyjew diffeomorphism*. Giving $T(T^*M)$ the tangent lift of the symplectic structure on T^*M , this Θ is a symplectic diffeomorphism to the canonical structure on $T^*(TM)$.

Anyone who meets these diffeomorphisms for the first time must surely suspect that they are related. For example, one would like (2) and (3) to be duals of (1). However, it is not at first clear what one might mean by this.

The best way to understand these maps is to extend them into a more general framework.

2.

In this section we consider those aspects of the constructions which do not depend upon the bracket of vector fields or the dual symplectic structure.

First consider T^2M . It has two vector bundle structures, the standard structure as the tangent bundle of TM , for which the projection is denoted p_{TM} , and what we call the *prolongation* structure, with projection $T(p_M)$ and operations which are obtained by applying the tangent functor T to the operations in TM . These two structures are represented in the diagram:

$$\begin{array}{ccc} & T(p_M) & \\ T^2M & \longrightarrow & TM \\ & \downarrow & \downarrow \\ p_{TM} & & \\ TM & \longrightarrow & M \end{array}$$

It is easier to keep the role of the two structures clear in one's mind if one replaces TM by an arbitrary vector bundle (E, q, M) :

$$(1) \quad \begin{array}{ccc} TE & \xrightarrow{T(q)} & TM \\ p_E \downarrow & & \downarrow \\ E & \longrightarrow & M \end{array}$$

Question: If we dualize $TE \rightarrow E$, is there a second structure which completes the square in

$$\begin{array}{ccc} T^*E & & ? \\ \downarrow & & \\ E & \longrightarrow & M \end{array}$$

To see the answer, consider the two short exact sequences associated with TE :

$$\begin{array}{c} q^!E \rightrightarrows TE \twoheadrightarrow q^!TM \\ p_M^! \rightrightarrows TE \twoheadrightarrow p_M^!E \end{array}$$

Here shrieks denote pullback bundles. The first sequence is a short exact sequence of vector bundles over base E and the surjection is the map induced by $T(q)$. The second sequence is over TM and the surjection is induced by p_E . The two surjections are actually the same map $TE \rightarrow E \times_M TM$, but are considered with respect to different bundle structures.

The kernel of the first surjection is the vertical bundle of q ; that is, the bundle of vectors tangent to the fibres of E . Such a vector is determined by its base point and by a vector in E itself to which it is parallel, and the injection $q^!E \rightarrow TE$ maps (e_1, e_2) to the vertical tangent vector with base point e_1 and parallel to e_2 .

The kernel of the second surjection consists of the tangent vectors to E along the zero section. Such a vector ξ may be written as $T(0)(x) + e$ where $x = T(q)(\xi) \in T_m(M)$ and $e \in E_m$ is identified with the vertical vector based at 0_m and parallel to e .

Thus the E which occurs in both kernels is precisely the set of tangent vectors to E which are vertical and based at points of the zero section; that is, it is $\ker(T(q)) \cap \ker(p_E)$.

Dualizing the first short exact sequence gives

$$q^!T^*M \rightrightarrows T^*E \twoheadrightarrow q^!E^*$$

and this suggests that the answer to the question is

$$(2) \quad \begin{array}{ccc} T^*E & \longrightarrow & E^* \\ \downarrow & & \downarrow \\ E & \longrightarrow & M \end{array}$$

That this is indeed so was proved in [9].

To find the dual of the other structure, $TE \longrightarrow TM$, apply T to the pairing $E \times_M E^* \longrightarrow \mathbb{R}$. This gives a pairing $TE \times_{TM} T(E^*) \longrightarrow \mathbb{R}$ which is again non-degenerate. So

$$(3) \quad (TE \longrightarrow TM)^* \cong (T(E^*) \longrightarrow TM)$$

and so dualizing the horizontal structure in (1) gives

$$(4) \quad \begin{array}{ccc} T(E^*) & \longrightarrow & TM \\ \downarrow & & \downarrow \\ E^* & \longrightarrow & M \end{array}$$

This is precisely the structure (1) with E replaced by E^* .

Now consider more general structures of this type.

A *double vector bundle* ([10]; see also [5, §1]) is a manifold D with two vector bundle structures, on bases A and B , each of which is a vector bundle on base M ,

$$\begin{array}{ccc} D & \xrightarrow{\tilde{q}_H} & B \\ \tilde{q}_V \downarrow & & \downarrow q_B \\ A & \xrightarrow{q_A} & M \end{array}$$

such that the structure maps of each vector bundle structure on D are vector bundle morphisms with respect to the other. Precisely: for the horizontal structure,

- the projection $\tilde{q}_H: D \longrightarrow B$ is a vector bundle morphism over the projection $q_A: A \longrightarrow M$;
- the zero section $B \longrightarrow D$ is a vector bundle morphism over the zero section $M \longrightarrow A$;
- the addition $D \times_B D \longrightarrow D$ is a vector bundle morphism over the addition $A \times_M A \longrightarrow A$;
- the scalar multiplication $D \times \mathbb{R} \longrightarrow D$ is a vector bundle morphism over the scalar multiplication $A \times \mathbb{R} \longrightarrow A$.

The first three conditions are precisely those which ensure that for four elements $d_1, d_2, d_3, d_4 \in D$ such that the LHS of

$$(d_1 \underset{H}{+} d_2) \underset{V}{+} (d_3 \underset{H}{+} d_4) = (d_1 \underset{V}{+} d_2) \underset{H}{+} (d_3 \underset{V}{+} d_4)$$

is defined, the RHS is defined and the two sides are equal. That (1) is indeed a double vector bundle is easily verified.

Now consider the intersection of the two kernels

$$C = \ker(\tilde{q}_H) \cap \ker(\tilde{q}_V).$$

A priori, there is no reason to expect this to have a natural algebraic structure. However, each structure on D induces a vector bundle structure on C with base M , and these two structures coincide. Pradines' terminology in [10] was *cœur*; we call it the *core* of D .

There is again a short exact sequence

$$q_A^! C \twoheadrightarrow D \twoheadrightarrow q_A^! B$$

of vector bundles over A and dualizing this gives

$$q_A^! B^* \twoheadrightarrow D^{*V} \twoheadrightarrow q_A^! C^*$$

and we get another double vector bundle

$$\begin{array}{ccc} D^{*V} & \longrightarrow & C^* \\ \downarrow & & \downarrow \\ A & \longrightarrow & M \end{array}$$

called the *vertical dual* of D . Likewise there is a *horizontal dual*

$$\begin{array}{ccc} D^{*H} & \longrightarrow & B \\ \downarrow & & \downarrow \\ C^* & \longrightarrow & M \end{array}$$

Theorem 1: ([6], [3]) The vector bundles $D^{*V} \rightarrow C^*$ and $D^{*H} \rightarrow C^*$ are themselves dual under a pairing which is canonical up to sign and which respects the double vector bundle structures.

To define the pairing, consider elements $\Phi \in D^{*V}$ and $\Psi \in D^{*H}$, shown with their various projections as follows:

$$\begin{array}{ccc} \Phi & \longmapsto & \kappa \\ \downarrow & & \downarrow \\ a & \longmapsto & m \end{array} \qquad \begin{array}{ccc} \Psi & \longmapsto & b \\ \downarrow & & \downarrow \\ \kappa & \longmapsto & m \end{array}$$

where $a \in A$, $b \in B$ and $\kappa \in C^*$. The given elements must of course lie over the same point of C^* . Now given a and b over the same $m \in M$, there exist elements

$d \in D$ of the form

$$\begin{array}{ccc} d & \longmapsto & b \\ \downarrow & & \downarrow \\ a & \longmapsto & m \end{array}$$

and we define

$$\langle \Phi, \Psi \rangle_{C^*} = \langle \Phi, d \rangle_A - \langle \Psi, d \rangle_B$$

where the subscripts indicate the bases of the pairings.

We could equally well reverse the order on the RHS. The statement in the theorem that this duality respects the double vector bundle structures means that the induced isomorphism from the dual of $D^{*V} \rightarrow C^*$ to $D^{*H} \rightarrow C^*$ is in fact an isomorphism of double vector bundles.

Example: Consider $D = TE$ as in (1). We observed that the two duals are $D^{*V} = T^*E$ as in (2) and $D^{*H} = T(E^*)$ as in (4). So Theorem 1 implies that there is a non-degenerate pairing of the two bundles

$$T^*E \rightarrow E^* \quad \text{and} \quad T(E^*) \rightarrow E^*.$$

To recognize what this is, write $A = E^*$. Then the statement is that the two bundles

$$T^*(A^*) \rightarrow A \quad \text{and} \quad T^*(A) \rightarrow A$$

are isomorphic as double vector bundles. That this is so was proved (by different methods) in [9]. To state the result precisely: Given a vector bundle $A \rightarrow M$, there is a canonical isomorphism of double vector bundles $R_A: T^*(A^*) \rightarrow T^*(A)$ over A and A^* , which is an anti-symplectomorphism and which induces the negative of the identity on the cores $T^*M \rightarrow T^*M$.

A proof of the next result is also in [9].

Proposition: For any manifold M , the composite

$$T^*(T^*M) \xrightarrow{\pi_{d\theta}^\#} T(T^*M) \xrightarrow{\Theta} T^*(TM)$$

is equal to R_{TM} .

The significance of this is that R is defined for arbitrary vector bundles, without reference to any structures of bracket type. Thus the Tulczyjew isomorphism could be intrinsically defined in terms of the symplectic structure on T^*M and the map R . (The definition in [12] uses the canonical involution on T^2M and the pairing which induces (3); see the example at the end of §3 below.)

Before we move on to consider the role of bracket structures in this context, it is interesting to observe that Theorem 1 is forced upon us by the simple pursuit of diagrams. Take an arbitrary double vector bundle and apply the tangent functor to get the triple structure in Figure 1(a). (The diagrams should be viewed as coming out of the page to the right.) Now dualize the bundle $TD \rightarrow D$. The rear square is precisely of the form (1), but when we dualize D we do so over B and so the dual is the horizontal dual. Likewise the left side face is of the form (1) but when we dualize D we do so over A , and so obtain the vertical dual. Now it is easy to guess that the eighth vertex must be C^* and one can then verify that Figure 1(b) is indeed a triple vector bundle. Using the isomorphisms R for the left and rear sides, we know that $T^*D \cong T^*(D^{*V})$ and $T^*D \cong T^*(D^{*H})$ as double vector bundles and




FIGURE 1.

we are forced to the conclusion that the top face must also be of the form (2), and that D^{*V} and D^{*H} must be duals.

3.

Bracket structures arise naturally from the differentiation of Lie groupoids. To proceed further, we thus need to consider general double Lie groupoids as in [5].

$$\begin{array}{ccc}
 S & \rightrightarrows & V \\
 \parallel & & \parallel \\
 H & \rightrightarrows & M
 \end{array}
 \tag{5}$$

Applying the Lie functor to the vertical structures gives the diagram on the left below [5]. That the Lie algebroid $A_V S$ of the vertical structure inherits a groupoid structure with base AV follows directly by observing that the Lie functor — like the tangent functor — preserves diagrams and pullbacks. The horizontal groupoid structures and the vertical Lie algebroid structures are compatible in the same way as in the definition of a double vector bundle above. Now applying the Lie functor a second time, to the horizontal structures, gives the diagram on the right [7]. This is a double Lie algebroid, but for the moment we will be concerned with it as a double vector bundle.

$$\begin{array}{ccc}
 A_V S & \rightrightarrows & AV \\
 \downarrow & & \downarrow \\
 H & \rightrightarrows & M
 \end{array}
 \tag{6}$$

$$\begin{array}{ccc}
 A_H(A_V S) & \longrightarrow & AV \\
 \downarrow & & \downarrow \\
 AH & \longrightarrow & M
 \end{array}$$

We could equally well apply the Lie functor first horizontally and then vertically, obtaining the following structures.

$$(7) \quad \begin{array}{ccc} A_H S & \longrightarrow & V \\ \Downarrow & & \Downarrow \\ A_H & \longrightarrow & M \end{array} \quad \begin{array}{ccc} A_V(A_H S) & \longrightarrow & A_V \\ \downarrow & & \downarrow \\ A_H & \longrightarrow & M \end{array}$$

The concept of core extends to double groupoids [2], [5] and in the case of S above yields a Lie groupoid K on base M , the Lie algebroid of which is the core of the two double vector bundles $A_V(A_H S)$ and $A_H(A_V S)$.

The Lie functor is a kind of generalized tangent functor, and both $A_V(A_H S)$ and $A_H(A_V S)$ are submanifolds of $T^2 S$. Further, the canonical involution J carries $A_V(A_H S)$ onto $A_H(A_V S)$ [7]. Denote this restricted map by j_S . This is an isomorphism of double vector bundles, as in Figure 2.

$$\begin{array}{ccccc} A_H(A_V S) & \longrightarrow & A_V & & \\ \downarrow & \searrow j_S & \downarrow & & \\ A_H & \longrightarrow & M & \xrightarrow{\quad} & A_V(A_H S) \longrightarrow A_V \\ & & & & \downarrow \qquad \downarrow \\ & & & & A_H \longrightarrow M \end{array}$$

FIGURE 2.

Dualizing j_S over A_H we obtain a map

$$j^{*V}: A_V^*(A_H S) \longrightarrow A_H^\bullet(A_V S).$$

The structure on the domain is the ordinary dual of the Lie algebroid of a Lie groupoid. On the target, however, the solidus indicates that we have taken the dual of a prolongation structure. As with the case

$$\left(\begin{array}{c} TE \\ \downarrow \\ TM \end{array} \right)^* \cong \left(\begin{array}{c} T(E^*) \\ \downarrow \\ TM \end{array} \right)$$

in (3), we can show that

$$\left(\begin{array}{c} A_H(A_V S) \\ \downarrow \\ AH \end{array} \right)^* \cong \left(\begin{array}{c} A(A_V^* S) \\ \downarrow \\ AH \end{array} \right)$$

where

$$(8) \quad \begin{array}{ccc} A_V^* S & \Longrightarrow & A^* K \\ \downarrow & & \downarrow \\ H & \Longrightarrow & M \end{array}$$

is a structure obtained by applying to $A_V S$ an extended form of the duality for double vector bundles (see [11] or [6]).

Denote the isomorphism $A(A_V^* S) \longrightarrow A_H^\bullet(A_V S)$ by I_V . We then define a generalization of the Tulczyjew isomorphism by

$$\Theta = I_V^{-1} \circ j^{*V} : A_V^*(A_H S) \longrightarrow A(A_V^* S)$$

(In [6] the inverse of this map is denoted j'^V .)

Example: To recover the classical case, consider the double groupoid

$$S = \begin{array}{ccc} M^4 & \Longrightarrow & M^2 \\ \Downarrow & & \Downarrow \\ M^2 & \Longrightarrow & M \end{array}$$

in which elements are quadruples of points, regarded as the corners of empty squares. Applying the Lie functor vertically, we obtain

$$A_V S = \begin{array}{ccc} TM \times TM & \Longrightarrow & TM \\ \downarrow & & \downarrow \\ M \times M & \Longrightarrow & M \end{array}$$

Here $TM \times TM$ is both the tangent Lie algebroid of $M \times M$ and the pair groupoid on base TM . The dualization process then produces

$$A_V^* S = \begin{array}{ccc} T^* M \times T^* M & \Longrightarrow & T^* M \\ \downarrow & & \downarrow \\ M \times M & \Longrightarrow & M \end{array}$$

in which again the horizontal groupoid structure is a pair groupoid. Applying the Lie functor to this horizontally we obtain

$$A(A_V^* S) = \begin{array}{ccc} T(T^* M) & \longrightarrow & T^* M \\ \downarrow & & \downarrow \\ TM & \longrightarrow & M \end{array}$$

The reader can similarly check that $A_V^*(A_H S) = T^*(TM)$. In this case I_V is precisely the isomorphism induced by the tangent pairing (3) of $T^2 M$ and $T(T^* M)$, and the definition of Θ in this case thus reduces exactly to that in [12].

4.

Finally we consider a general form of the map $T^*(T^*M) \longrightarrow T(T^*M)$ obtained from the symplectic structure on a cotangent bundle.

Theorem 2: [6] Given any double Lie groupoid (5) with core groupoid $K \rightrightarrows M$, there is a structure of double Lie groupoid on T^*S which makes it a symplectic double groupoid

$$\begin{array}{ccc} T^*S & \rightrightarrows & A_V^*S \\ \Downarrow & & \Downarrow \\ A_H^*S & \rightrightarrows & A^*K \end{array}$$

and which induces on A_V^*S , A_H^*S and A^*K their standard Poisson structures. In particular, $A_H^*S \rightrightarrows A^*K$ and $A_V^*S \rightrightarrows A^*K$ are Poisson groupoids in duality.

The statement that two Poisson groupoids are in duality means that the Lie algebroid of each, constructed from the Lie groupoid structure, is isomorphic (or anti-isomorphic) to the Lie algebroid dual of the other with the Lie algebroid structure which it acquires from the Poisson groupoid structure [13]. (We will not consider how to arrange signs here; there are various possibilities.) Thus Theorem 2 asserts the existence of Lie algebroid isomorphisms

$$\mathcal{D}_H: A^*(A_H^*S) \longrightarrow A(A_V^*S), \quad \mathcal{D}_V: A^*(A_V^*S) \longrightarrow A(A_H^*S).$$

Following through the Example at the end of §3, the reader can verify that these generalize the map $\pi_{d\theta}^\#$. Whereas $\pi_{d\theta}^\#$ arises from the symplectic structure on T^*M , the maps $\mathcal{D}_H$ and $\mathcal{D}_V$ arise from the Poisson structures on the duals of Lie algebroids.

In order to express the relationship between these maps $\mathcal{D}$ and the Tulczyjew diffeomorphism in the general case, we need one more ingredient. Consider the structure A_H^*S from (7). Applying the duality process mentioned in connection with (8), we obtain the structure on the left below. Applying the Lie functor vertically we obtain the double vector bundle on the right below.

$$(9) \quad \begin{array}{ccc} A_H^*S & \longrightarrow & V \\ \Downarrow & & \Downarrow \\ A^*K & \longrightarrow & M \end{array} \quad \begin{array}{ccc} A(A_H^*S) & \longrightarrow & AV \\ \downarrow & & \downarrow \\ A^*K & \longrightarrow & M \end{array}$$

Now consider the vertical dual of the double vector bundle $A_V(A_H^*S)$ in (7). By an extension of the methods used in §2, one can define a non-degenerate pairing of the double vector bundles $A(A_H^*S)$ and $A_V^*(A_H^*S)$ which induces an isomorphism of double vector bundles

$$R_H: A_V^*(A_H^*S) \longrightarrow A^*(A_H^*S).$$

In the case where S is actually a double vector bundle, this reduces to the R of §2.

Theorem 3:[6] For a double Lie groupoid S as in (5), the following diagram of diffeomorphisms commutes.

$$\begin{array}{ccc}
A(A_V^*S) & \xrightarrow{\mathcal{D}_H^{-1}} & A^*(A_H^*S) \\
I_V \downarrow & \searrow \Theta & \downarrow R_H \\
A^\bullet(A_HS) & \xrightarrow{(j^{*V})^{-1}} & A^*(A_HS)
\end{array}$$

In particular, it follows that Θ is a Poisson diffeomorphism, since both $\mathcal{D}_H$ and R_H are anti-Poisson [6].

In conclusion, it is worthwhile noticing the different degrees of development required by the various canonical diffeomorphisms which we have discussed. The map $R_A: T^*(A^*) \rightarrow T^*(A)$ is the simplest, requiring only a single ordinary vector bundle. The Tulczyjew diffeomorphism requires the notion of double Lie groupoid, and of its associated double Lie algebroid, but involves only the simpler aspects of that construction — a dualization of the map j as a morphism of double vector bundles and the prolongation of a pairing which exists on the level of the intermediate structures A_VS . The maps $\mathcal{D}_V$ and $\mathcal{D}_H$, on the other hand, require a detailed consideration of the structure of the symplectic double groupoid associated with a double Lie groupoid. As was shown in [6] and [8], this structure is the key to the general notion of double Lie algebroid.

References

- [1] R. Abraham and J. Marsden. *Foundations of Mechanics*. Addison-Wesley, second edition, 1985.
- [2] R. Brown and K. C. H. Mackenzie. Determination of a double Lie groupoid by its core diagram. *J. Pure Appl. Algebra*, 80(3):237–272, 1992.
- [3] Katarzyna Konieczna and Paweł Urbański. Double vector bundles and duality. *Arch. Math. (Brno)*, 35(1):59–95, 1999.
- [4] Zhang-Ju Liu, Alan Weinstein, and Ping Xu. Manin triples for Lie bialgebroids. *J. Differential Geom.*, 45:547–574, 1997.
- [5] K. C. H. Mackenzie. Double Lie algebroids and second-order geometry, I. *Adv. Math.*, 94(2):180–239, 1992.
- [6] K. C. H. Mackenzie. On symplectic double groupoids and the duality of Poisson groupoids. *Internat. J. Math.*, 10:435–456, 1999.
- [7] K. C. H. Mackenzie. Double Lie algebroids and second-order geometry, II. *Adv. Math.*, 154:46–75, 2000.
- [8] K. C. H. Mackenzie. Notions of double for Lie algebroids. math.dg/0011212, 2000. 30 pages.
- [9] K. C. H. Mackenzie and Ping Xu. Lie bialgebroids and Poisson groupoids. *Duke Math. J.*, 73(2):415–452, 1994.
- [10] J. Pradines. Fibrés vectoriels doubles et calcul des jets non holonomes. Notes polycopiées, Amiens, 1974.
- [11] J. Pradines. Quotients de groupoïdes différentiables. *C. R. Acad. Sci. Paris Sér. I Math.*, 303:817–820, 1986.
- [12] W. M. Tulczyjew. *Geometric formulation of physical theories*, volume 11 of *Monographs and Textbooks in Physical Science*. Bibliopolis, Naples, 1989.

- [13] A. Weinstein. Coisotropic calculus and Poisson groupoids. *J. Math. Soc. Japan*, 40:705–727, 1988.

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF SHEFFIELD, SHEFFIELD, S3 7RH,
UNITED KINGDOM

E-mail address: `K.Mackenzie@sheffield.ac.uk`